

# Research Paper: The Anti-Parkinsonism Effects of $K_{ATP}$ Channel Blockade in the 6-Hydroxydopamine-Induced Animal Model: The Role of Oxidative Stress



Hossein Piri<sup>1</sup>, Hashem Haghdoust-Yazdi<sup>1\*</sup>, Negin Fraidouni<sup>2</sup>, Tahereh Dargahi<sup>2</sup>, Mohamadhosein Yaghoubidoust<sup>2</sup>, Abbas Azadmehr<sup>3</sup>

1. Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.

2. Student Research Committee, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran.

3. Department of Immunology, School of Medicine, Babol University of Medical Sciences, Babol, Iran.



**Citation:** Piri, H., Haghdoust Yazdi, H., Fraidoun, N., Dargahi, T., Yaghoubidoust, M., & Azadmehr, A. (2017). The Anti-Parkinsonism Effects of  $K_{ATP}$  Channel Blockade in the 6-Hydroxydopamine-Induced Animal Model: The Role of Oxidative Stress. *Basic and Clinical Neuroscience*, 8(3), 183-192. <https://doi.org/10.18869/nirp.bcn.8.3.183>

**doi:** <https://doi.org/10.18869/nirp.bcn.8.3.183>

## Article info:

Received: 06 September 2016

First Revision: 05 December 2016

Accepted: 20 December 2016

## ABSTRACT

**Introduction:** Studies suggest that ATP-sensitive potassium ( $K_{ATP}$ ) channels are a potential pharmacotherapeutic target for neuroprotection in neurodegenerative diseases. The current study aimed at evaluating the effect of pretreatment with glibenclamide (Glib) and B vitamins supplement on the severity of behavioral symptoms in 6-hydroxydopamine (OHDA)-induced Parkinsonism. Also malondialdehyde (MDA) concentration was measured in the blood and brain suspensions to find probable neuroprotective mechanism of Glib.

**Methods:** The 6-OHDA was injected into striatum of rats by stereotaxic surgery. Treatment with Glib and B vitamins was started before the surgery and continued up to 3 weeks after that. Development and severity of Parkinsonism were evaluated by conventional behavioral tests. MDA values were measured spectrophotometrically using thiobarbituric acid and MDA standard curve.

**Results:** Pretreatments with Glib, at both doses of 1 and 5 mg/kg or B vitamins significantly ameliorated severity of the behavioral symptoms. Pretreatment with a combination of Glib and B vitamins was more effective than pretreatment with Glib or B vitamins alone. Also, pretreatment with B vitamins, Glib, or a combination of them reduced MDA concentration in the brain suspensions. Decrease in MDA concentration in the group of rats that received a combination of B vitamins and Glib was more prominent than that of the Glib groups.

**Conclusion:** As severity of the behavioral symptoms in the 6-OHDA-induced Parkinsonism reflects the degree of the lesion in Substantia Nigra (SN) dopaminergic neurons, it is suggested that Glib pretreatment has neuroprotective effect against 6-OHDA-induced neurotoxicity. The current study data also showed that this effect may be mediated by antioxidant effect of Glib.

## Key Words:

Parkinson disease,  
6-Hydroxydopamine,  
Glibenclamide,  
B vitamins,  
Malondialdehyde

\* Corresponding Author:

**Hashem Haghdoust-Yazdi, PhD**

**Address:** Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.

**Tel:** +98 (281) 3366001

**E-mail:** hhaghdoust@yahoo.com